



MM3D-Fluorogenic Probe for Real-Time Measurement of ENGase Activity



MM3D is hydrolyzed by endo- β -*N*-acetylglucosaminidase (ENGase) to generate fluorescence. MM3D is a useful tool for real-time measurement of the enzymatic activity of ENGase.

Principle



Endo- β -*N*-acetylglucosaminidase (ENGase) first hydrolyzes the chitobiose moiety of *N*-glycan. MM3D, a *N*-glycan which features both a fluorescent group (*N*-methyl anthraniloyl, MANT) and a quencher group (dinitrophenyl, DNP) in the structure. Cleavage of MM3D by ENGase removes the self-quenching effect, leading to an increase in fluorescence intensity. MM3D has shown to be useful for high throughput analysis with plate readers, and in the search for new types of ENGase or inhibitor of ENGase. These inhibitors can be served as therapeutic agents for NGLY1-deficiency, a very rare genetic disease.


*Note: MM3D is not hydrolyzed by Endo-H which belongs to GH Family 18, and therefore is not suitable for measurement of Endo-H activity.

- References**
- 1) N. Ishii, C. Sunaga, K. Sano, C. Huang, K. Iino, Y. Matsuzaki, T. Suzuki, I. Matsuo, *ChemBioChem* **2018**, *19*, 660.
 - 2) H. Fujihira, Y. Masahara-Negishi, M. Tamura, C. Huang, Y. Harada, S. Wakana, D. Takakura, N. Kawasaki, N. Taniguchi, G. Kondoh, T. Yamashita, Y. Funakoshi, T. Suzuki, *PLoS Genet.* **2017**, *13*, e1006696.
 - 3) C. Huang, Y. Harada, A. Hosomi, Y. Masahara-Negishi, J. Seino, H. Fujihira, Y. Funakoshi, T. Suzuki, N. Dohmae, T. Suzuki, *Proc. Natl. Acad. Sci. U.S.A.* **2015**, *112*, 1398.

This product was commercialized by collaboration with Prof. Ichiro Matsuo.

MM3D-Fluorogenic Probe for Real-Time Measurement of ENGase Activity

Measurement of Endo-M [A1651] activity using MM3D¹⁾



(Data are provided by Prof. Ichiro Matsuo.)

Comparing the fluorescence intensity of MM3D with that of the hydrolyzed product, the fluorescence of MANT is decreased by 98% due to the quenching effect of the DNP group (a). However, since the cleaved product of MM3D is highly sensitive, the concentration of MM3D is sufficient in 0.5 μ M to measure Endo-M [A1651] activity (b). Next, addition of DMSO to hinder the enzymatic reaction suppresses enzyme activity in a concentration-dependent manner (c). These results indicate that MM3D is suitable for screening for potential inhibitors of ENGase.

Related Product

endo- β -N-Acetylglucosaminidase (= Endo-M)
Recombinant: from *Mucor hiemalis* expressed in *Candida boidinii*

1 vial [A1651]

For further information please refer to our website at www.TCIchemicals.com.

glycoscience



Ordering and Customer Service

TCI AMERICA

Tel : 800-423-8616 / 503-283-1681
Fax : 888-520-1075 / 503-283-1987
E-mail : Sales-US@TCIchemicals.com

TCI EUROPE N.V.

Tel : +32 (0)3 735 07 00
Fax : +32 (0)3 735 07 01
E-mail : Sales-EU@TCIchemicals.com

TCI Deutschland GmbH

Tel : +49 (0)6196 64053-00
Fax : +49 (0)6196 64053-01
E-mail : Sales-DE@TCIchemicals.com

Tokyo Chemical Industry UK Ltd.

Tel : +44 (0)1865 78 45 60
E-mail : Sales-UK@TCIchemicals.com

TCI Chemicals (India) Pvt. Ltd.

Tel : 1800 425 7889 / 044-2262 0909
Fax : 044-2262 8902
E-mail : Sales-IN@TCIchemicals.com

梯希爱(上海)化成工业发展有限公司

Tel : 800-988-0390 / 021-67121386
Fax : 021-6712-1385
E-mail : Sales-CN@TCIchemicals.com

TOKYO CHEMICAL INDUSTRY CO., LTD.

Tel : +81 (0)3-5640-8878
E-mail : globalbusiness@TCIchemicals.com

Availability, price or specification of the listed products are subject to change without prior notice. Reproduction forbidden without the prior written consent of Tokyo Chemical Industry Co., Ltd.