

N-Mesityl Substituted Chiral Triazolium Salts: Opening a New World of N-Heterocyclic Carbene Catalysis

Pei-Chen Chiang and Jeffrey W. Bode



Laboratorium für Organische Chemie
ETH-Zürich
Zürich 8093, Switzerland
bode@org.chem.ethz.ch

1. Introduction

The dimerization of aldehydes to form benzoin (**Scheme 1**) is one of the oldest known organic reactions, first reported in 1832 by Liebig.¹ The original cyanide catalyzed benzoin reaction was long appreciated for its simplicity and mechanistic intricacies but offered few useful synthetic applications.² The discovery by Ukai³ in 1943 that the essential co-factor thiamine, or vitamin B1, catalyzes the benzoin reaction provided a key clue to understanding a broad range of thiamine-dependent enzymes and reinvigorated interest in this class of reactions.⁴ The use of ylides derived from thiamine, which are now classified as N-heterocyclic carbenes,⁵ led to both new reactions, such as the Stetter addition of aldehydes to electron-deficient olefins,⁶ and new catalysts types inspired by the chemistry of the thiazolium salt at the heart of thiamine.⁷ Key to the success of thiamine and its relative is the catalytic generation of an acyl anion equivalent.⁸ Typical procedures for the preparation of this reactive species would require harsh conditions and

intricate protecting group strategies. With thiamine catalysis, this reactive species can be generated under aqueous and nearly neutral conditions.

By the year 2003, modern, enantioselective versions of the benzoin and certain Stetter reactions had emerged, driven by the design and preparation of new chiral azolium salts. Pioneering work by Knight and Leeper,⁹ Enders,¹⁰ and Rovis¹¹ had converged on the use of chiral, bicyclic triazolium salts as the ideal platform for N-heterocyclic carbene catalysts capable of inducing highly enantioselective reactions including intermolecular homo-benzoin reactions,¹² intramolecular Stetter reactions,¹³ and intramolecular aldehyde–ketone crossed benzoin reactions.¹⁴ These methods are now well established and widely employed for the enantioselective synthesis of complex molecules including bioactive natural products.¹⁵



Scheme 1. Benzoin reaction with benzaldehydes.


3. Catalyst Synthesis and Counterion Effects

Our catalyst design borrows its chiral elements on the outstanding work of Knight and Leeper,⁹ who first reported chiral triazolium salts derived from 1,2-amino alcohols, and of Rovis,¹¹ who first employed the readily available 1,2-aminoindanol for chiral azolium salts. In both cases, high quality *N*-mesityl-hydrazine hydrochloride is needed. While this is commercially available the cost is prohibitive for large scale syntheses. We have therefore devised a reliable procedure for the large scale preparation of this hydrazine salt from the corresponding aniline. Although yields are not high (36–40% yield), it can be executed on a 1 mol scale with inexpensive starting materials and without the need for chromatography or distillation.


The preparation of the chiral catalysts begins by following the work of Rovis, who has prepared aminoindanol derived triazoliums bearing different aromatic substituents.¹³ Lactam

1 is converted to the imidate **2** with trimethyloxonium tetrafluoroborate and condensation with *N*-mesityl hydrazine affords **3** (Scheme 4). The most challenging step is the final ring closing, which fails or gives low yields under the previously developed conditions. Our modified procedure uses lower temperatures and HCl or HClO₄ to promote the ring closure. The resulting chloride or perchlorate salts are readily isolated and stored.¹⁸

In all cases examined to date, the counterion plays no role in the reaction outcome. Thus catalysts of type **A** or **B** bearing chloride, perchlorate or tetrafluoroborate counterions give identical yields and enantioselectivities provided that at least a catalytic amount of base is used in the reactions (Scheme 5).¹⁹ The only discernable difference is that the chloride catalysts are somewhat more hygroscopic but this does not generally affect the reaction outcome.



Scheme 4. Synthesis of chiral triazolium catalyst *ent-B*.




Scheme 5. In the presence of bases, there is no significant effect of the azolium counterions. (DIPEA = *N,N*-diisopropylethylamine, DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene)


4. Hetero-Diels–Alder Reactions

The catalytic generation of ester enolate equivalents under mild, simple conditions that allow for enantioselective reactions has been a long held goal of synthetic organic chemistry. In 2006, we disclosed the use of *N*-mesityl substituted triazolium salts for the catalytic generation of ester enolate equivalents from α -halo- and α,β -unsaturated aldehydes.


Racemic α -halo aldehydes undergo highly enantioselective annulations with electron deficient oxo-dienes under remarkably mild and simple reaction conditions (**Scheme 6**).²⁰ Due to the high reactivity of these substrates, only 0.5 mol% of chiral triazolium salt **B** is needed for excellent yields and enantio-selectivities. The racemic α -halo aldehydes undergo epimerization in the reaction conditions rendering this reaction



Scheme 6. Diels–Alder reactions of α -chloroaldehydes with oxodienes.



Scheme 7. Diels–Alder reactions with α -chloroaldehyde bisulfite salts.



Scheme 8. Diels–Alder reactions with α,β -unsaturated aldehydes. (NMM = *N*-methylmorpholine, DMAP = 4-(dimethylamino)pyridine)

an enantio-convergent process. The scope of this process is outstanding, with either aromatic or aliphatic substituents tolerated by both reaction partners.


The α -halo aldehyde starting materials are readily prepared in a single step from the corresponding aldehydes. They have, however, limited shelf life. We have therefore developed conditions for isolating and storing them as their bisulfite adducts and developed biphasic reaction conditions for the use of these convenient starting materials directly in the enantioselective annulation reactions (**Scheme 7**).²¹ Similarly high yields, enantioselectivities, and substrate scope are observed even under aqueous conditions. Importantly, this procedure works well with the commercially available bisulfite adduct of chloroacetaldehyde, making possible enantioselective acetate additions without the need to prepare the toxic and potentially explosive α -chloroaldehyde.

The identical reactive intermediates can be generated from α,β -unsaturated aldehydes and we have recently reported a method for highly enantioselective annulations from both aromatic and aliphatic substituted enals (**Scheme 8**).²² Under these conditions, high catalyst loadings are needed due to the


fact that only a trace amount of the active carbene catalysts is generated by the weak bases. This procedure also expands the scope of the annulations to include α -hydroxy- and α -aminoenones that give products bearing valuable synthetic handles for further transformations. Scheidt has also applied these chiral catalysts to an intra-molecular variant of this reaction.²³

The heterodiene can also be extended to nitrogen analogues using either α -chloroaldehydes or commercially available electron-deficient enals as the enolate precursors (**Scheme 9**).¹⁷ This procedure affords *cis*-disubstituted dihydropyridinones in good yields and with outstanding enantioselectivities.

Recently, several other groups have used our catalyst and conditions for even more complex hetero-Diels–Alder reactions. An excellent example is that of Kobayashi and coworkers, who reported the formation of amine-substituted products by reactions of α -haloaldehydes and vinylogous amides (**Scheme 10**).²⁴



Scheme 9. Azadiene Diels–Alder reactions using α,β -unsaturated aldehydes.



Scheme 10. Enantioselective synthesis of dihydropyranones.

5. Catalytic Generation of Homoenolate Equivalents




The nature of the azolium catalyst employed in the annulation reactions plays an important role in the type of reactive intermediate generated. α,β -Unsaturated aldehydes can access either the homoenolate or enolate pathways. In general, the enolate pathway is favored by the use of weak bases (DMAP or NMM) and triazolium catalysts, while stronger bases (DBU) and imidazolium-derived N-heterocyclic carbenes prefer the homoenolate equivalent route. Reactions that proceed via the homoenolate pathway generate five-membered ring products such as γ -lactams, γ -lactones, or cyclopentane derivatives. For example, the same substrates shown in **Scheme 11** give either γ -lactams or dihydropyridinones depending on the catalyst type.¹⁷

In general, imidazolium-derived catalysts are superior to triazolium-derived variants for γ -lactone²⁵ and γ -lactam²⁶ formation, with commercially available IMesCl (cat. C, CAS

Nr. 141556-45-8) as the best achiral catalysts for many of these transformations (**Scheme 12**).

For certain highly reactive electrophiles, *N*-mesityl substituted triazolium salts are also active. The actual reaction mechanisms may be more complex in these cases, but the products are those derived from formal homoenolate generation and addition. Several such examples are listed below.

Saccharine derived-cyclic ketimines are unexpectedly good electrophiles in NHC-catalyzed annulations with enals using achiral triazolium salt **A** (**Scheme 13**).²⁷ These reactions demonstrate broad scope in both the enal and ketimine reaction partner and afford unique polycyclic products in excellent yield and with low catalysts loading. Although chiral triazolium salt **B** is also an excellent catalyst for this reaction, enantioselectivities are relatively modest. This reflects the known difficulty of effecting highly enantioselective reactions that proceed via the homoenolate pathway.






When α,β -unsaturated *N*-tosyl ketimines are used as substrates, bicyclic β -lactams, rather than γ -lactams are formed (**Scheme 14**).²⁸ This reaction is most easily thought of as a conjugate addition of the homoenolate derived from the α,β -unsaturated aldehyde followed by ring-closing reactions, although the actual reaction mechanism probably involves a different pathway. This remarkable cascade process affords enantiopure β -lactams in excellent yields and stereoselectivities.

A similar reaction for the formation of enantiopure cyclopentyl β -lactones is also possible with the same catalyst

and unsaturated aldehydes (**Scheme 15**).²⁹ In this case, the use of α' -hydroxyenones is essential and provides a useful synthetic handle for the elaboration of the products.

With enones bearing an aromatic group, the initial β -lactone products undergo spontaneous decarboxylation to afford disubstituted-cyclopentenones (**Scheme 16**). The overall reaction allows the combination of an enal and a simple enone to give enantioenriched cyclopentenones in a single step and under mild, simple reaction conditions.³⁰




6. Catalytic Generation of Activated Carboxylates

In all of the reactions noted above, catalyst turnover is affected by the formation of an acyl azolium species that undergoes a lactonization or lactamization reaction. These acyl azolium intermediates therefore serve as catalytically generated activated carboxylates and can be used for esterification and amidation reactions. By choosing the appropriate substrates, *N*-heterocyclic carbene catalysts can effect both esterifications and amidations under catalytic conditions that do not require coupling reagents or produce chemical waste.


The first reports on the catalytic generation of acyl azoliums for esterification reactions utilized simpler azolium salts. In 2004, our group reported the diastereoselective opening of α,β -epoxyaldehydes with simple thiazolium-derived ylides (**Scheme 17**).³¹ The *N*-mesityl substituted triazolium salt **A** also catalyzes this reaction and is in many cases the superior

reagent. Rovis, also in 2004, reported the generation of acyl azoliums from α -halo aldehydes using *N*-phenyl substituted triazolium salt **E** (**Scheme 18**).³² Rovis has also described an impressive example of enantioselective protonation using a chiral *N*-pentafluorophenyl substituted triazolium salt.³³


Although a number of azolium precatalysts can be used for redox esterification reactions of α -functionalized aldehydes, the *N*-mesityl or *N*-pentafluorophenyl substituted triazolium salts usually give superior results. A rapidly growing body of literature on these reactions has established that almost any aldehyde with an α -leaving group or reversible functionality serves as a substrate for NHC-catalyzed redox esterifications. The *N*-mesityl substituted catalyst is particularly suited for redox reaction of α,β -unsaturated aldehydes (**Scheme 19**).¹⁶ Other applications include the opening of enantioenriched formyl cyclopropanes to generate esters³⁴ or dihydropyranones reported by You and coworkers.³⁵




Scheme 17. Formation of esters from α,β -epoxyaldehydes.



Scheme 18. Formation of esters from α -bromoaldehydes.



Scheme 19. Redox reactions promoted by catalyst **A**.




Scheme 20. Redox amidations.

A curious finding of both our group and several others is the reluctance of catalytically generated acyl azoliums to undergo reactions with amines to form amides. This property can be exploited for the chemoselective acylation of alcohols in the presence of amines. The origin of this unusual reactivity lies in the unique properties of the acyl azoliums themselves, which do not readily react with amines to form amide products.³⁶ Successful NHC-catalyzed amidations do occur with the addition of suitable co-catalysts. In our hands, we have found that the addition of imidazole or similar triazoles make catalytic amidations of α -functionalized aldehydes possible, via the intermediacy of an acyl imidazole that reacts with the amine (**Scheme 20**).³⁷ Similar chemistry using HOAt as the co-catalytic acylating agent was reported simultaneously by Rovis.³⁸


7. Catalytic Annulations by NHC-Catalyzed Claisen Rearrangements

The combination of a chiral N-heterocyclic carbene catalysts and an ynals should lead to the formation of an α,β -unsaturated acyl azolium that may act as an electrophile, representing a completely different activation mode than the acyl anion, homoenolate, ester enolate, or activated carboxylate equivalents discussed so far. Using the identical chiral catalyst, we succeeded in identifying conditions for the selective generation of such intermediates and their use in annulation reactions with enolic substrates, such as kojic acid derivatives (**Scheme 21**).³⁹

Our initial work focused on kojic acids as substrates due to their known synthetic utility, but similar conditions give annulation products from pyruvates in excellent yields and selectivities. Recently reported chemistry from other groups



Scheme 21. Enantioselective annulations of ynals and kojic acids.



Scheme 22. Mechanism of Claisen rearrangements.

using 1,3-dicarbonyls as nucleophiles are likely to proceed via a similar mechanism and should be amenable to enantioselective catalysis with chiral catalyst **B** or its relatives.⁴⁰ The reactions may also be conducted by starting from the α,β -unsaturated aldehyde and a stoichiometric oxidant with no effect to the observed enantioselectivities.³⁹

The mechanism of this reaction is also of considerable interest. Although we initially believed the α,β -unsaturated acyl azolium **4** would be an excellent Michael acceptor, we have been unsuccessful in adding any nucleophiles other than enolates or MeOH/water. This led us to extensively investigate the detailed mechanism revealing an NHC-catalyzed Coates-Claisen rearrangement as the key step. This mechanistic postulate rationalizes the exceptional enantioselectivities observed in this reaction (**Scheme 22**).


8. α' -Hydroxyenones as Enal Surrogates

The majority of the reaction described in this review use α,β -unsaturated aldehydes as starting materials. Although not obscure compounds, the vast majority of substrates require multiple steps for their preparation. This is particularly true of the highly desired heteroaromatic substrates for which the methods of enal synthesis are complicated by the presence of basic functionality.


In order to improve the scope of the NHC-catalyzed reactions and expand the range of products that may be prepared with these methods, we have sought readily prepared, bench stable surrogates for aldehydes reaction partners. For ester enolate equivalents, the α -chloroaldehyde bisulfite adducts are ideal,²¹ but these cannot be used for reactions involving homoenolate equivalents.

We have therefore developed α' -hydroxyenones **6** as readily prepared surrogates for α,β -unsaturated aldehydes.⁴¹ These substrates can be prepared on a multigram scale from commercially available ketone **5** and aromatic aldehydes (**Scheme 23**).

In the presence of an N-heterocyclic carbene, the α' -hydroxyenones undergo identical reactions as the corresponding α,β -unsaturated aldehydes. In this case, the key reactive intermediates are formed by a retro-benzoin reaction of the initial NHC-ketone adduct (**Scheme 24**).⁴²



Scheme 23. Base catalyzed aldol reactions of α' -hydroxyenones synthesis.





Scheme 24. Generations of Breslow intermediate from α' -hydroxyenones and enals.

These substrates are particularly well suited for the synthesis of cyclopentenes and lactams, with selected examples shown in **Scheme 25**. Due to the increased steric demands of the ketone substrates, these reactions work best with achiral triazolium precatalyst **A**. We therefore highly recommend these conditions for the preparation of racemic mixtures and the preparation of small libraries. Should the enantiopure compounds be needed, they can be prepared with chiral catalysts **B** and *ent-B* from the α,β -unsaturated aldehyde.

The α -hydroxyenones are also outstanding substrates for NHC-catalyzed amidations in combination with 1,2,4-triazole as a co-catalyst.⁴³ Unlike most amidation conditions that require excess amounts of coupling reagents and generate a large amount of waste, these conditions affect catalytic amidations with no products other than acetone and only substoichiometric quantities of reagents. Selected examples are shown in **Scheme 26**.

9. Conclusions

In just four years since the initial report, *N*-mesityl substituted chiral triazolium salts have proven to be one of the most versatile and selective catalysts known. In addition to the remarkable range of products they may be used to construct, they are notable for their high stability, mild reaction conditions (20–40 °C), and tolerance to air and water. A single chiral catalysts work well for nearly all of the reactions reported to date. For a few NHC-catalyzed reactions for which **B** and *ent-B* give inferior enantioselectivities, a number of new chiral scaffolds are emerging. Notable, all of these structures maintain the essential *N*-mesityl triazolium core that has proved to be necessary for both reactivity and selectivity in these new generations of *N*-heterocyclic carbene catalyzed reactions. The commercial availability of these catalysts should encourage the discovery of even more new reactions and applications of the complex, enantiopure products they afford.



References

- 1 F. Wöhler, J. von Liebig, *Annalen der Pharmacie* **1832**, *3*, 249–282.
- 2 A. Lapworth, *J. Chem. Soc., Trans.* **1903**, *83*, 995–1005.
- 3 T. Ukai, R. Tanaka, S. Dokawa, *J. Pharm. Soc. Jpn.* **1943**, *63*, 296–300.
- 4 General review: R. Kluger, K. Tittmann, *Chem. Rev.* **2008**, *108*, 1797–1833.
- 5 H. W. Wanzlick, E. Schikora, *Angew. Chem.* **1960**, *72*, 494.
- 6 General review: H. Stetter, H. Kuhlmann, in *The Catalyzed Nucleophilic Addition of Aldehydes to Electrophilic Double Bonds*, John Wiley & Sons, **1991**, Vol. 40, pp 407–496.
- 7 General review: D. Enders, A. A. Narine, *J. Org. Chem.* **2008**, *73*, 7857–7870.
- 8 R. Breslow, *J. Am. Chem. Soc.* **1958**, *80*, 3719–3726.
- 9 (a) R. L. Knight, F. J. Leeper, *Tetrahedron Lett.* **1997**, *38*, 3611–3614. (b) R. L. Knight, F. J. Leeper, *J. Chem. Soc., Perkin Trans. 1* **1998**, 1891–1893.
- 10 (a) D. Enders, K. Breuer, J. H. Teles, *Helv. Chim. Acta* **1996**, *79*, 1217–1221. (b) D. Enders, K. Breuer, J. Runsink, J. H. Teles, *Helv. Chim. Acta* **1996**, *79*, 1899–1902.
- 11 M. S. Kerr, J. Read de Alaniz, T. Rovis, *J. Org. Chem.* **2005**, *70*, 5725–5728.
- 12 D. Enders, U. Kallfass, *Angew. Chem. Int. Ed.* **2002**, *41*, 1743–1745.
- 13 M. S. Kerr, J. Read de Alaniz, T. Rovis, *J. Am. Chem. Soc.* **2002**, *124*, 10298–10299.
- 14 H. Takikawa, Y. Hachisu, J. W. Bode, K. Suzuki, *Angew. Chem. Int. Ed.* **2006**, *45*, 3492–3494.
- 15 General review: (a) J. Moore, T. Rovis, Carbene Catalysts. In *Topics in Current Chemistry*, Springer Berlin, Heidelberg, **2009**, Vol. 291, pp 77–144. (b) F. Glorius, K. Hirano, Nucleophilic Carbenes as Organocatalysts. In *Ernst Schering Found.*, Springer Berlin, Heidelberg, **2008**, Vol. 2, pp 159–181.
- 16 S. S. Sohn, J. W. Bode, *Org. Lett.* **2005**, *7*, 3873–3876.
- 17 M. He, J. R. Struble, J. W. Bode, *J. Am. Chem. Soc.* **2006**, *128*, 8418–8420.
- 18 J. R. Struble, J. W. Bode, *Org. Synth.* **2010**, *87*, 362–376.
- 19 J. R. Struble, J. Kaeobamrung, J. W. Bode, *Org. Lett.* **2008**, *10*, 957–960.
- 20 M. He, G. J. Uc, J. W. Bode, *J. Am. Chem. Soc.* **2006**, *128*, 15088–15089.
- 21 M. He, B. J. Beahm, J. W. Bode, *Org. Lett.* **2008**, *10*, 3817–3820.
- 22 J. Kaeobamrung, M. C. Kozlowski, J. W. Bode, *Proc. Nat. Acad. Sci. U.S.A.* **2010**, *107*, 20661–20665.
- 23 E. M. Phillips, M. Wadamoto, A. Chan, K. A. Scheidt, *Angew. Chem. Int. Ed.* **2007**, *46*, 3107–3110.
- 24 S. Kobayashi, T. Kinoshita, H. Uehara, T. Sudo, I. Ryu, *Org. Lett.* **2009**, *11*, 3934–3937.
- 25 S. S. Sohn, E. L. Rosen, J. W. Bode, *J. Am. Chem. Soc.* **2004**, *126*, 14370–14371.
- 26 M. He, J. W. Bode, *Org. Lett.* **2005**, *7*, 3131–3134.
- 27 M. Rommel, T. Fukuzumi, J. W. Bode, *J. Am. Chem. Soc.* **2008**, *130*, 17266–17267.
- 28 M. He, J. W. Bode, *J. Am. Chem. Soc.* **2008**, *130*, 418–419.
- 29 J. Kaeobamrung, J. W. Bode, *Org. Lett.* **2009**, *11*, 677–680.
- 30 P.-C. Chiang, J. Kaeobamrung, J. W. Bode, *J. Am. Chem. Soc.* **2007**, *129*, 3520–3521.
- 31 K. Y. Chow, J. W. Bode, *J. Am. Chem. Soc.* **2004**, *126*, 8126–8127.
- 32 N. T. Reynolds, J. Read de Alaniz, T. Rovis, *J. Am. Chem. Soc.* **2004**, *126*, 9518–9519.
- 33 H. U. Vora, T. Rovis, *J. Am. Chem. Soc.* **2010**, *132*, 2860–2861.
- 34 S. S. Sohn, J. W. Bode, *Angew. Chem. Int. Ed.* **2006**, *45*, 6021–6024.
- 35 G.-Q. Li, L.-X. Dai, S.-L. You, *Org. Lett.* **2009**, *11*, 1623–1625.
- 36 (a) K. Daigo, L. Reed, *J. Am. Chem. Soc.* **1962**, *84*, 659–662. (b) T. C. Owen, A. Richards, *J. Am. Chem. Soc.* **1987**, *109*, 2520–2521. (c) T. C. Owen, J. N. Harris, *J. Am. Chem. Soc.* **1990**, *112*, 6136–6137.
- 37 J. W. Bode, S. S. Sohn, *J. Am. Chem. Soc.* **2007**, *129*, 13798–13799.
- 38 H. U. Vora, T. Rovis, *J. Am. Chem. Soc.* **2007**, *129*, 13796–13797.
- 39 J. Kaeobamrung, J. Mahatthananchai, P. Zheng, J. W. Bode, *J. Am. Chem. Soc.* **2010**, *132*, 8810–8812.
- 40 (a) S. J. Ryan, L. Candish, D. W. Lupton, *J. Am. Chem. Soc.* **2009**, *131*, 14176–14177. (b) Z. Q. Zhu, J. C. Xiao, *Adv. Synth. Catal.* **2010**, *352*, 2455–2458. (c) S. De Sarkar, A. Studer, *Angew. Chem. Int. Ed.* **2010**, *49*, 9266–9269.
- 41 (a) G. G. King, J. V. Karabinos, *J. Chem. Eng. Data* **1968**, *13*, 565–566. (b) C. Palomo, M. Oiarbide, J. M. García, A. González, E. Arceo, *J. Am. Chem. Soc.* **2003**, *125*, 13942–13943. (c) M. Reiter, H. Turner, R. Mills-Webb, V. Gouverneur, *J. Org. Chem.* **2005**, *70*, 8478–8485.
- 42 P.-C. Chiang, M. Rommel, J. W. Bode, *J. Am. Chem. Soc.* **2009**, *131*, 8714–8718.
- 43 P.-C. Chiang, Y. Kim, J. W. Bode, *Chem. Commun.* **2009**, 4566–4568.

(Received October 2010)

Introduction of the authors :

Pei-Chen Chiang

Graduate student, Department of Chemistry, University of Pennsylvania

[Education and employment] 2004.1 M.S., National Chiao-Tung University, Taiwan (Supervisor: Prof. Wen-Sheng Chung); 2004-2005 Research Assistant, National Chiao-Tung University, Taiwan (Supervisor: Prof. Eric W. G. Diau); 2006.6-present: Ph.D. student, University of Pennsylvania, USA

[Specialties] Chemistry, Organic Synthesis, Asymmetric Catalysis Chemistry

Jeffrey W. Bode


Professor, Laboratory of Organic Chemistry, Department of Chemistry and Applied Biosciences, Swiss Federal Institute of Technology (ETH)-Zürich, Switzerland

[Education and employment] 1996 B.S., Trinity University, San Antonio, Texas, USA (Supervisor: Prof. Michael P. Doyle); 2001 Dok. Nat. Sci., Swiss Federal Institute of Technology (ETH)-Zürich, Switzerland (Supervisor: Prof. Erick M. Carreira); 2001-2003 Postdoctoral Research Fellow, Tokyo Institute of Technology, Tokyo, Japan (Supervisor: Keisuke Suzuki);


2003-2007 Assistant Professor of Chemistry and Biochemistry, University of California, Santa Barbara, USA; 2007-2009 Associate Professor of Chemistry, University of Pennsylvania, USA; 2010-present Professor of Synthetic Organic Chemistry, Swiss Federal Institute of Technology (ETH)-Zürich, Switzerland
[Specialties] Chemistry, Organic Synthesis, Asymmetric Catalysis Chemistry, Peptide Chemistry

Contribution Related Compounds


Bode Catalysts



6,7-Dihydro-2-(2,4,6-trimethylphenyl)-
5H-pyrrolo[2,1-c]-1,2,4-triazolium
Perchlorate
100mg [D3982]



Bode Catalyst 2
Monohydrate
100mg [D3983]



Bode Catalyst 1
Monohydrate
100mg [D3984]